

Physics 704/804 Electromagnetic Theory II

G. A. Krafft

Jefferson Lab

Jefferson Lab Professor of Physics

Old Dominion University

Applied to Spherical Harmonics



$$L_+ Y_{lm} = \sqrt{(l-m)(l+m+1)} Y_{l, m+1}$$

$$L_- Y_{lm} = \sqrt{(l+m)(l-m+1)} Y_{l, m-1}$$

$$L_z Y_{lm} = m Y_{lm}$$

$$L^2 \vec{L} = \vec{L} L^2$$

$$\vec{L} \times \vec{L} = i\vec{L}$$

$$L_j \nabla^2 = \nabla^2 L_j$$

Vector Spherical Harmonics



$$\vec{X}_{lm} = \frac{1}{\sqrt{l(l+1)}} \vec{L} Y_{lm}(\theta, \phi)$$

$$\int \vec{X}_{l'm'}^* \cdot \vec{X}_{lm} d\Omega = \delta_{ll'} \delta_{mm'}$$

Did exercises in Chapter 12.11 of Arfken (3rd Edition) and obtained a lot of clarity

$$\text{Show } \vec{X}_{lm} = \frac{1}{\sqrt{l(l+1)}} \vec{L} Y_{lm} =$$

$$\hat{\theta} \left\{ \frac{-m}{\sqrt{l(l+1)} \sin \theta} Y_{lm} \right\} + \hat{\phi} \left\{ \frac{-i}{\sqrt{l(l+1)}} \frac{\partial Y_{lm}}{\partial \theta} \right\}$$

Verify the orthogonality relation

$$\int \vec{X}_{l'm'}^* \cdot \vec{X}_{lm} d\Omega = \delta_{ll'} \delta_{mm'}$$

Verify the sum rule

$$\sum_{m=-l}^l \left| \vec{X}_{lm}(\theta, \phi) \right|^2 = \frac{2l+1}{4\pi}$$

Show power for pure multipole formula from last time:

$$\begin{aligned} \frac{dP(l, m)}{d\Omega} &= \frac{Z_0}{2k^2} |a(l, m)|^2 \left| \vec{X}_{lm} \right|^2 \\ &= \frac{Z_0}{2k^2 l(l+1)} |a(l, m)|^2 \left\{ \begin{aligned} &\frac{1}{2}(l-m)(l+m+1) |Y_{l, m+1}|^2 \\ &+ \frac{1}{2}(l+m)(l-m+1) |Y_{l, m-1}|^2 + m^2 |Y_{l, m}|^2 \end{aligned} \right\} \end{aligned}$$

Multipole Expansion



$$\left(\nabla^2 + k^2\right) \vec{H} = 0 \quad \vec{\nabla} \cdot \vec{H} = 0 \quad \vec{E} = \frac{iZ_0}{k} \vec{\nabla} \times \vec{H}$$

$$\left(\nabla^2 + k^2\right) \vec{E} = 0 \quad \vec{\nabla} \cdot \vec{E} = 0 \quad \vec{H} = -\frac{i}{kZ_0} \vec{\nabla} \times \vec{E}$$

$$\nabla^2 (\vec{r} \cdot \vec{A}) = \vec{r} \cdot (\nabla^2 \vec{A}) + 2\vec{\nabla} \cdot \vec{A}$$

$$\left(\nabla^2 + k^2\right) (\vec{r} \cdot \vec{H}) = 0 \quad \left(\nabla^2 + k^2\right) (\vec{r} \cdot \vec{E}) = 0$$

Magnetic Multipole Field (TE)



$$\left(\vec{r} \cdot \vec{H}_{lm}^M\right) = \frac{l(l+1)}{k} g_l(kr) Y_{lm}(\theta, \phi)$$

$$\left(\vec{r} \cdot \vec{E}_{lm}^M\right) = 0$$

$$Z_0 k \vec{r} \cdot \vec{H} = \vec{L} \cdot \vec{E}$$

$$\vec{L} \cdot \vec{E}_{lm}^M(r, \theta, \phi) = l(l+1) Z_0 g_l(r) Y_{lm}(\theta, \phi)$$

$$\vec{E}_{lm}^M = Z_0 g_l(r) \vec{L} Y_{lm}(\theta, \phi)$$

$$\vec{H}_{lm}^M = -\frac{i}{kZ_0} \vec{\nabla} \times \vec{E}_{lm}^M$$

Electric Multipole (TM)

$$\left(\vec{r} \cdot \vec{E}_{lm}^E\right) = -Z_0 \frac{l(l+1)}{k} f_l(kr) Y_{lm}(\theta, \phi)$$

$$\left(\vec{r} \cdot \vec{H}_{lm}^E\right) = 0$$

$$\vec{H}_{lm}^E = f_l(r) \vec{L} Y_{lm}(\theta, \phi)$$

$$\vec{E}_{lm}^E = \frac{iZ_0}{k} \vec{\nabla} \times \vec{H}_{lm}^E$$

normal mode expansions

$$\vec{X}_{lm} = \frac{1}{\sqrt{l(l+1)}} \vec{L} Y_{lm}(\theta, \phi)$$

$$\int \vec{X}_{l'm'}^* \cdot \vec{X}_{lm} d\Omega = \delta_{ll'} \delta_{mm'}$$

$$\int \vec{X}_{l'm'}^* \cdot \left(\vec{r} \times \vec{X}_{lm}\right) d\Omega = 0$$

General Solution



$$\vec{H} = \sum_{lm} \left[a_E(l, m) f_l(kr) \vec{X}_{lm} - \frac{i}{k} a_M(l, m) \vec{\nabla} \times g_l(kr) \vec{X}_{lm} \right]$$

$$\vec{E} = \sum_{lm} \left[\frac{i}{k} a_E(l, m) \vec{\nabla} \times f_l(kr) \vec{X}_{lm} + a_M(l, m) g_l(kr) \vec{X}_{lm} \right]$$

$$a_M(l, m) g_l(kr) = \frac{k}{\sqrt{l(l+1)}} \int Y_{lm}^* \vec{r} \cdot \vec{H} d\Omega$$

$$Z_0 a_E(l, m) g_l(kr) = - \frac{k}{\sqrt{l(l+1)}} \int Y_{lm}^* \vec{r} \cdot \vec{E} d\Omega$$

\therefore knowledge at 2 different radii sufficient for complete knowledge of solution

Energy



$$\vec{H}_l = \sum_m a_E(l, m) \vec{X}_{lm} h_l^1(kr) e^{-i\omega t}$$

$$\vec{E}_l = \frac{i}{k} Z_0 \vec{\nabla} \times \vec{H}_l$$

$$u = \frac{\epsilon_0}{4} (\vec{E} \cdot \vec{E}^* + Z_0^2 \vec{H} \cdot \vec{H}^*) \approx \frac{\mu_0}{2} \vec{H} \cdot \vec{H}^*$$

in radiation zone

$$dU = \frac{\mu_0 dr}{2k^2} \sum_{mm'} a_E^*(l, m) a_E(l, m') \int \vec{X}_{lm'}^* \cdot \vec{X}_{lm} d\Omega$$

$$\frac{dU}{dr} = \frac{\mu_0}{2k^2} \sum_{mm'} |a_E(l, m)|^2$$

incoherent sum over multipoles

Angular Momentum



$$\vec{m} = \frac{1}{2c^2} \text{Re} \left[\vec{r} \times \left(\vec{E} \times \vec{H}^* \right) \right]$$

$$\vec{m} = \frac{\mu_0}{2\omega} \text{Re} \left[\vec{H}^* \left(\vec{L} \cdot \vec{H} \right) \right]$$

$$\frac{d\vec{M}}{dr} = \frac{\mu_0}{2\omega k^2} \text{Re} \sum_{mm'} a_E^*(l, m) a_E(l, m') \int Y_{lm'}^* \vec{L} Y_{lm} d\Omega$$

$$\frac{dM_z}{dr} = \frac{\mu_0}{2\omega k^2} \sum_m m |a_E(l, m)|^2$$

$$\frac{dM_z}{dr} = \frac{m}{\omega} \frac{dU}{dr}$$

photon interpretation: photon takes away
 $m\hbar$ units of angular momentum