

Accelerator Physics Particle Acceleration

G. A. Krafft Old Dominion University Jefferson Lab Lecture 8







Statistical Treatments of Beams

- Distribution Functions Defined
 - Statistical Averaging
 - Examples
- Kinetic Equations
 - Liouville Theorem
 - Vlasov Theory
 - Collision Corrections
- Self-consistent Fields
- Collective Effects
 - KV Equation
 - Landau Damping
- Beam-Beam Effect





Beam rms Emittance



Treat the beam as a statistical ensemble as in Statistical Mechanics. Define the distribution of particles within the beam statistically. Define single particle distribution function

 $\psi(x,x'),$

where $\psi(x,x')dxdx'$ is the number of particles in [x,x+dx] and [x',x'+dx'], and statistical averaging as in Statistical Mechanics, e. g.

$$\left\langle q\right\rangle \equiv \int q\left(x, x'\right)\psi\left(x, x'\right)dxdx' / N$$
$$\left\langle q^{2}\right\rangle \equiv \int q^{2}\left(x, x'\right)\psi\left(x, x'\right)dxdx' / N$$
$$\vdots$$



Thomas Jefferson National Accelerator Facility



Closest *rms* **Fit Ellipses**



For zero-centered distributions, i.e., distributions that have zero average value for x and x'

$$\varepsilon_{rms} \equiv \sqrt{\left\langle x^2 \right\rangle \left\langle x'^2 \right\rangle - \left\langle xx' \right\rangle^2}$$



$$\gamma = \frac{\langle x'^2 \rangle}{\varepsilon_{rms}} = \frac{\sigma_{x'}^2}{\varepsilon_{rms}}$$



Thomas Jefferson National Accelerator Facility



Case: Uniformly Filled Ellipse



$$\psi(x,x') = \frac{1}{\pi\varepsilon} \Theta\left(1 - \frac{\gamma x^2 + 2\alpha x x' + \beta x'^2}{\varepsilon}\right)$$

 Θ here is the Heavyside step function, 1 for positive values of its argument and zero for negative values of its argument

$$\sigma_x^2 = \langle x^2 \rangle = \frac{\varepsilon \beta}{4}$$
$$\langle xx' \rangle = -\frac{\varepsilon \alpha}{4}$$
$$\sigma_{x'}^2 = \langle x'^2 \rangle = \frac{\varepsilon}{4\beta} (1 + \alpha^2)$$



Thomas Jefferson National Accelerator Facility



Graduate Accelerator Physics Fall 2015

 $\therefore \mathcal{E}_{rms} = \frac{c}{\Lambda}$

Dynamics? Start with Liouville Thm

Generalization of the Area Theorem of Linear Optics. Simple Statement: For a dynamical system that may be described by a conserved energy function (Hamiltonian), the relevant phase space volume is conserved by the flow, *even if the forces are non-linear*. Start with some simple geometry!



Case: Uniformly Filled Ellipse



$$\psi(x,x') = \frac{1}{\pi\varepsilon} \Theta\left(1 - \frac{\gamma x^2 + 2\alpha x x' + \beta x'^2}{\varepsilon}\right)$$

 Θ here is the Heavyside step function, 1 for positive values of its argument and zero for negative values of its argument

$$\sigma_x^2 = \langle x^2 \rangle = \frac{\varepsilon \beta}{4}$$
$$\langle xx' \rangle = -\frac{\varepsilon \alpha}{4}$$
$$\sigma_{x'}^2 = \langle x'^2 \rangle = \frac{\varepsilon}{4\beta} (1 + \alpha^2)$$



Thomas Jefferson National Accelerator Facility



Graduate Accelerator Physics Fall 2015

 $\therefore \mathcal{E}_{rms} = \frac{c}{\Lambda}$

Dynamics? Start with Liouville Thm

Generalization of the Area Theorem of Linear Optics. Simple Statement: For a dynamical system that may be described by a conserved energy function (Hamiltonian), the relevant phase space volume is conserved by the flow, *even if the forces are non-linear*. Start with some simple geometry!



Liouville Theorem

$$\begin{aligned} &(q_{0}, p_{0}) \longrightarrow \left(q_{0} + \frac{\partial H}{\partial p}(q_{0}, p_{0})\Delta t + \dots, p_{0} - \frac{\partial H}{\partial q}(q_{0}, p_{0})\Delta t + \dots\right) \\ &(q_{0} + \Delta q, p_{0}) \longrightarrow \left(q_{0} + \Delta q + \frac{\partial H}{\partial p}(q_{0} + \Delta q, p_{0})\Delta t + \dots, p_{0} - \frac{\partial H}{\partial q}(q_{0} + \Delta q, p_{0})\Delta t + \dots\right) \\ &(q_{0}, p_{0} + \Delta p) \longrightarrow \left(q_{0} + \frac{\partial H}{\partial p}(q_{0}, p_{0} + \Delta p)\Delta t + \dots, p_{0} + \Delta p - \frac{\partial H}{\partial q}(q_{0}, p_{0} + \Delta p)\Delta t + \dots\right) \\ &(q_{0} + \Delta q, p_{0} + \Delta p) \longrightarrow \left(q_{0} + \Delta q + \frac{\partial H}{\partial p}(q_{0} + \Delta q, p_{0} + \Delta p)\Delta t + \dots, p_{0} + \Delta p - \frac{\partial H}{\partial q}(q_{0}, p_{0} + \Delta p)\Delta t + \dots\right) \\ &(q_{0} + \Delta q, p_{0} + \Delta p) \longrightarrow \left(q_{0} + \Delta q + \frac{\partial H}{\partial p}(q_{0} + \Delta q, p_{0} + \Delta p)\Delta t + \dots\right) \\ &(q_{0} + \Delta q, p_{0} + \Delta p) \longrightarrow \left(q_{0} + \Delta q - \frac{\partial H}{\partial q}(q_{0} + \Delta q, p_{0} + \Delta p)\Delta t + \dots\right) \\ &\therefore \text{Area } \Delta_{1} = \frac{1}{2} \det \left| \frac{\Delta q}{\partial p} \left(q_{0} + \Delta q, p_{0}\right) - \frac{\partial H}{\partial p}(q_{0}, p_{0})\right]\Delta t \qquad - \left[\frac{\partial H}{\partial q}(q_{0} + \Delta q, p_{0}) - \frac{\partial H}{\partial q}(q_{0}, p_{0})\right]\Delta t \\ & \left[\frac{\partial H}{\partial p}(q_{0}, p_{0} + \Delta p) - \frac{\partial H}{\partial p}(q_{0}, p_{0})\right]\Delta t \qquad \Delta p - \left[\frac{\partial H}{\partial q}(q_{0}, p_{0} + \Delta p) - \frac{\partial H}{\partial q}(q_{0}, p_{0})\right]\Delta t \\ & \underset{M \to 0}{\longrightarrow} \frac{1}{2} \left(\Delta q \Delta p\right) \left[1 + \left\{\frac{\partial^{2} H}{\partial q \partial p}(q_{0}, p_{0}) - \frac{\partial^{2} H}{\partial p \partial q}(q_{0}, p_{0})\right\}\Delta t\right] = \frac{(\Delta q \Delta p)}{2} \end{aligned}$$



Thomas Jefferson National Accelerator Facility





Likewise

$$\operatorname{Area} \Delta_{2} = \frac{1}{2} \operatorname{det} \begin{vmatrix} \Delta q + \left[\frac{\partial H}{\partial p} (q_{0} + \Delta q, p_{0}) - \frac{\partial H}{\partial p} (q_{0} + \Delta q, p_{0} + \Delta p) \right] \Delta t & - \left[\frac{\partial H}{\partial q} (q_{0} + \Delta q, p_{0}) - \frac{\partial H}{\partial q} (q_{0} + \Delta q, p_{0} + \Delta p) \right] \Delta t \\ \left[\frac{\partial H}{\partial p} (q_{0}, p_{0} + \Delta p) - \frac{\partial H}{\partial p} (q_{0} + \Delta q, p_{0} + \Delta p) \right] \Delta t & \Delta p - \left[\frac{\partial H}{\partial q} (q_{0}, p_{0} + \Delta p) - \frac{\partial H}{\partial q} (q_{0} + \Delta q, p_{0} + \Delta p) \right] \Delta t \end{vmatrix}$$

$$\xrightarrow{\Delta t \to 0} \frac{1}{2} (\Delta q \Delta p) \left[1 + \left\{ \frac{\partial^{2} H}{\partial q \partial p} (q_{0} + \Delta p, p_{0} + \Delta p) - \frac{\partial^{2} H}{\partial p \partial q} (q_{0} + \Delta p, p_{0} + \Delta p) - \frac{\partial^{2} H}{\partial p \partial q} (q_{0} + \Delta p, p_{0} + \Delta p) \right\} \Delta t \right] = \frac{(\Delta q \Delta p)}{2}$$

Because the starting point is arbitrary, phase space area is conserved at each location in phase space. In three dimensions, the full 6-D phase volume is conserved by essentially the same argument, as is the sum of the projected areas in each individual projected phase spaces (the so-called third Poincare and first Poincare invariants, respectively). Defeat it by adding non-Hamiltonian (dissipative!) terms later.





Phase Space

• Plot of dynamical system "state" with coordinate along abscissa and momentum along the ordinate





Thomas Jefferson National Accelerator Facility



Liouville Theorem

- ODU
- Area in phase space is preserved when the dynamics is Hamiltonian





Thomas Jefferson National Accelerator Facility



1D Proof





Thomas Jefferson National Accelerator Facility



$$\frac{dV}{dt} = \int_{0}^{L} \left[-p_x(s) \frac{d}{ds} \frac{\partial H}{\partial p_x}(x(s), p_x(s)) + \frac{\partial H}{\partial x}(x(s), p_x(s)) \frac{dx(s)}{ds} \right] ds$$
$$= \int_{0}^{L} \left[\frac{dp_x(s)}{ds} \frac{\partial H}{\partial p_x}(x(s), p_x(s)) + \frac{\partial H}{\partial x}(x(s), p_x(s)) \frac{dx(s)}{ds} \right] ds$$

(why is boundary term of integration by parts zero?)

$$= \oint_{c(s,t)} \left[\frac{\partial H}{\partial p_x} dp_x + \frac{\partial H}{\partial x} dx \right]$$

By Green's Thm.

=0 when the differential is an exact differential

i.e.,
$$\frac{\partial}{\partial x} \left(\frac{\partial H}{\partial p_x} \right) = \frac{\partial}{\partial p_x} \left(\frac{\partial H}{\partial x} \right)$$
,

in other words always

(note the integrand above is really dH, so H is a "potential") for phase space!!!



Thomas Jefferson National Accelerator Facility



3D Poincare Invariants

- e 6D phase ODU
- In a three dimensional Hamiltonian motion, the 6D phase space volume is conserved (also called Liouville's Thm.) $V_{6D} = \int dp_x dp_y dp_z dx dy dz$

Additionally, the sum
$$V_6$$
 of the projected volumes (Poincare invariants) are conserved

$$\int_{\text{proj}(V_2)} dp_x dx + \int_{\text{proj}(V_2)} dp_y dy + \int_{\text{proj}(V_2)} dp_z dz$$

$$\int_{\text{proj}(V_4)} dp_y dp_z dy dz + \int_{\text{proj}(V_4)} dp_z dp_x dz dx + \int_{\text{proj}(V_4)} dp_x dp_y dx dy$$

Emittance (phase space area) exchange based on this idea

• More complicated to prove, but are true because, as in 1D $\partial^2 H$ $\partial^2 H$

$$\overline{\partial q_i \partial p_i} = \overline{\partial p_i \partial q_i}$$



Thomas Jefferson National Accelerator Facility





$\gamma \text{ is a loop in 6D phase space}$ $\gamma(t) = \left(\vec{p}(s,t), \vec{q}(s,t)\right)$ $\frac{d}{dt} \left[\oint_{\gamma(t)} \sum_{i=1}^{3} p_i dx_i \right] = \int_{0}^{L} \sum_{i=1}^{3} \left[-p_i(s) \frac{d}{ds} \frac{\partial H}{\partial p_i}(\vec{x}(s), \vec{p}(s)) + \frac{\partial H}{\partial x_i}(\vec{x}(s), \vec{p}(s)) \frac{dx_i(s)}{ds} \right]$ $= \int_{0}^{L} \sum_{i=1}^{3} \left[\frac{dp_i(s)}{ds} \frac{\partial H}{\partial p_i}(\vec{x}(s), \vec{p}(s)) + \frac{\partial H}{\partial x_i}(\vec{x}(s), \vec{p}(s)) \frac{dx_i(s)}{ds} \right] = \oint_{\gamma(t)} dH = 0$

for any surface in 6D phase space V_2 , with $\gamma = \partial V_2$

$$\oint_{\partial V_2} \sum_{i=1}^3 p_i dx_i = \int_{V_2} \sum_{i=1}^3 dp_i dx_i = \sum_{i=1}^3 \int_{\text{proj}(V_2)} dp_i dx_i$$

$$\left(\sum_{i=1}^{3} dp_{i} dx_{i}\right)^{2} = dp_{y} dp_{z} dy dz + dp_{z} dp_{x} dz dx + dp_{x} dp_{y} dx dy$$

$$\left(\sum_{i=1}^{3} dp_i dx_i\right)^3 = dp_x dp_y dp_z dx dy dz$$



Thomas Jefferson National Accelerator Facility



Vlasov Equation



By interpretation of ψ as the single particle distribution function, and because the individual particles in the distribution are assumed to not cross the boundaries of the phase space volumes (collisions neglected!), ψ must evolve so that

$$\frac{d\psi}{dt} = 0 \qquad \text{as the distibution evolves}$$

$$\frac{d\psi}{dt} = \lim_{\delta t \to 0} \frac{\psi(t + \delta t, \vec{q}(t + \delta t), \vec{p}(t + \delta t)) - \psi(t, \vec{q}(t), \vec{p}(t))}{\delta t} = 0$$
where the equation for ANY (this is what makes
it hard to solve in general!) individual orbits
through phase space is given by $\vec{q}(t), \vec{p}(t)$

$$\therefore \frac{\partial \psi}{\partial t} + \frac{d\vec{q}}{dt} \frac{\partial \psi}{\partial \vec{q}} + \frac{d\vec{p}}{dt} \frac{\partial \psi}{\partial \vec{p}} = 0$$



Thomas Jefferson National Accelerator Facility



Conservation of Probability



 $N(t) = \int \psi(t; \vec{q}, \vec{p}) d^3 \vec{x} d^3 \vec{p}$ is a conserved quantity

continuity equation for ψ is

$$\frac{\partial \psi}{\partial t} + \nabla_{\vec{q}} \left(\dot{\vec{q}} \psi \right) + \nabla_{\vec{p}} \left(\dot{\vec{p}} \psi \right) = 0$$

$$\therefore \frac{\partial \psi}{\partial t} + \frac{d\vec{q}}{dt} \frac{\partial \psi}{\partial \vec{q}} + \frac{d\vec{p}}{dt} \frac{\partial \psi}{\partial \vec{p}} + \psi \left[\nabla_{\vec{q}} \frac{\partial H}{\partial \vec{p}} - \nabla_{\vec{p}} \frac{\partial H}{\partial \vec{q}} \right] = 0$$

for the Hamiltonian system



Thomas Jefferson National Accelerator Facility



Jean's Theorem

The independent variable in the Vlasov equation is often changed to the variable *s*. In this case the Vlasov equation is

$$\frac{\partial \psi}{\partial s} + \frac{d\vec{q}}{ds}\frac{\partial \psi}{\partial \vec{q}} + \frac{d\vec{p}}{ds}\frac{\partial \psi}{\partial \vec{p}} = 0$$

The equilibrium Vlasov problem, $\partial \psi / \partial t = 0$, is solved by any function of the constants of the motion. This result is called Jean's theorem, and is the starting point for instability analysis as the "unperturbed problem".

If $\psi = f(A, B, C, \dots)$, where A, B, C, \dots are constants of the motion

$$\frac{d\vec{x}}{dt}\frac{\partial\psi}{\partial\vec{x}} + \frac{d\vec{p}}{dt}\frac{\partial\psi}{\partial\vec{p}} = \frac{\partial f}{\partial A}\left(\frac{d\vec{x}}{dt}\frac{\partial A}{\partial\vec{x}} + \frac{d\vec{p}}{dt}\frac{\partial A}{\partial\vec{p}}\right) + \frac{\partial f}{\partial B}\left(\frac{d\vec{x}}{dt}\frac{\partial B}{\partial\vec{x}} + \frac{d\vec{p}}{dt}\frac{\partial B}{\partial\vec{p}}\right) + \frac{\partial f}{\partial C}\left(\frac{d\vec{x}}{dt}\frac{\partial C}{\partial\vec{x}} + \frac{d\vec{p}}{dt}\frac{\partial C}{\partial\vec{p}}\right) + \dots = \frac{\partial f}{\partial A}\frac{dA}{dt} + \frac{\partial f}{\partial B}\frac{dB}{dt} + \frac{\partial f}{\partial C}\frac{dC}{dt} + \dots = 0$$



Thomas Jefferson National Accelerator Facility



Examples



1-D Harmonic oscillator Hamiltonian. Bi-Maxwellian distribution is a stationary distribution

$$\psi = \frac{1}{2\pi} \exp\left(-H/kT\right) = \frac{m\omega}{2\pi kT} \exp\left(-p_x^2/2mkT\right) \exp\left(-mx^2\omega^2/2kT\right),$$

As is any other function of the Hamiltonian. Contours of constant ψ line up with contours of constant *H*

2 D transverse Gaussians, including focusing structure in ring

$$\psi(s; x, x'; y, y') \propto \exp\left(-\left(\gamma_x(s)x^2 + 2\alpha_x(s)xx' + \beta_x(s)x'^2\right)/\varepsilon_x\right) \\ \times \exp\left(-\left(\gamma_y(s)y^2 + 2\alpha_y(s)yy' + \beta_y(s)y'^2\right)/\varepsilon_y\right)$$

Contours of constant ψ line up with contours of constant Courant-Snyder invariant. Stationary as particles move on ellipses!



Thomas Jefferson National Accelerator Facility



Solution by Characteristics

More subtle: a solution to the full Vlasov equation may be obtained from the distribution function at some the initial condition, provided the particle orbits may be found unambiguously from the initial conditions throughout phase space. Example: 1-D harmonic oscillator Hamiltonian.

$$\begin{pmatrix} x(t) \\ x'(t) \end{pmatrix} = \begin{pmatrix} \cos \omega (t-t_0) & \sin \omega (t-t_0) / \omega \\ -\omega \sin \omega (t-t_0) & \cos \omega (t-t_0) \end{pmatrix} \begin{pmatrix} x(t_0) \\ x'(t_0) \end{pmatrix} \rightarrow \begin{pmatrix} x_0 \\ x'_0 \end{pmatrix} = \begin{pmatrix} \cos \omega (t-t_0) & -\sin \omega (t-t_0) / \omega \\ \omega \sin \omega (t-t_0) & \cos \omega (t-t_0) \end{pmatrix} \begin{pmatrix} x \\ x' \end{pmatrix} \\ \psi (x, x'; t = t_0) = f_0 (x, x')$$
Let $\psi (x, x'; t) = f_0 (\cos \omega (t-t_0) x - \sin \omega (t-t_0) x' / \omega, \omega \sin \omega (t-t_0) x + \cos \omega (t-t_0) x')$

$$\frac{\partial \psi}{\partial t} = \frac{\partial f_0}{\partial x} \frac{dx(t; x, x')}{dt} + \frac{\partial f_0}{\partial x'} \frac{dx'(t; x, x')}{dt}$$

$$= \frac{\partial f}{\partial x} \left[-\omega \sin \omega (t-t_0) x - \cos \omega (t-t_0) x' \right] + \frac{\partial f}{\partial x'} \left[\omega^2 \cos \omega (t-t_0) x - \omega \sin \omega (t-t_0) x' \right]$$

$$\frac{dx}{dt} \frac{\partial \psi}{\partial x} = x' \left[\frac{\partial f}{\partial x} \cos \omega (t-t_0) + \frac{\partial f}{\partial x'} \omega \sin \omega (t-t_0) \right]$$



Thomas Jefferson National Accelerator Facility Graduate Accelerator Physics Fall 2015



Breathing Mode





The particle envelope "breaths" at twice the revolution frequency!



Thomas Jefferson National Accelerator Facility



Sacherer Theory



Assume beam is acted on by a linear focusing force plus additional linear or non-linear forces

$$x'' + k_x^2 x - F_x = 0$$

$$y'' + k_y^2 x - F_y = 0$$

For space charge example we'll see

$$F_{x(y)} = \frac{qE_{x(y)}\left(1-\beta^2\right)}{\gamma mc^2\beta^2} = \frac{qE_{x(y)}}{\gamma^3 mc^2\beta^2}$$

Now

$$\langle xx'' \rangle + k_x^2 \langle x^2 \rangle - \langle F_x x \rangle = 0 \langle yy'' \rangle + k_y^2 \langle y^2 \rangle - \langle F_y y \rangle = 0$$

Assume distributions zero-centered and let

$$\tilde{x}^2 = \langle x^2 \rangle \quad \tilde{x}'^2 = \langle x'^2 \rangle \qquad \tilde{y}^2 = \langle y^2 \rangle \quad \tilde{y}'^2 = \langle y'^2 \rangle$$



Thomas Jefferson National Accelerator Facility







Thomas Jefferson National Accelerator Facility



rms Emittance Conserved

$$\left(\left\langle x^2 \right\rangle \left\langle x'^2 \right\rangle - \left\langle xx' \right\rangle^2 \right)'$$

$$= \left\langle x^2 \right\rangle' \left\langle x'^2 \right\rangle + \left\langle x^2 \right\rangle \left\langle x'^2 \right\rangle' - 2 \left\langle xx' \right\rangle \left\langle xx' \right\rangle'$$

$$= 2 \left\langle xx' \right\rangle \left\langle x'^2 \right\rangle + 2 \left\langle x^2 \right\rangle \left\langle x'x'' \right\rangle - 2 \left\langle xx' \right\rangle \left(\left\langle x'^2 \right\rangle - k_x^2 \left\langle x^2 \right\rangle + \left\langle F_x x \right\rangle \right)$$

$$= 2 \left\langle x^2 \right\rangle \left(-k_x^2 \left\langle x'x \right\rangle + \left\langle F_x x' \right\rangle \right) + 2 \left\langle xx' \right\rangle \left(k_x^2 \left\langle x^2 \right\rangle - \left\langle F_x x \right\rangle \right)$$

$$= 2 \left\langle x^2 \right\rangle \left\langle F_x x' \right\rangle - 2 \left\langle xx' \right\rangle \left\langle F_x x \right\rangle$$

For linear forces derivative vanishes and *rms* emittance conserved. Emittance growth implies *non-linear forces*.





Space Charge and Collective Effects

- Collective Effects
 - Brillouin Flow
 - Self-consistent Field
 - KV Equation
 - Bennet Pinch
 - Landau Damping





Simple Problem



- How to account for interactions between particles
- Approach 1: Coulomb sums
 - Use Coulomb's Law to calculate the interaction between each particle in beam
 - Unfavorable N^2 in calculation but perhaps most realistic
 - more and more realistic as computers get better
- Approach 2: Calculate EM field using ME
 - Need procedure to define charge and current densities
 - Track particles in resulting field





Uniform Beam Example

• Assume beam density is uniform and axi-symmetric going into magnetic field



Electric Field

$$\frac{1}{r}\frac{\partial}{\partial r}rE_r = \frac{qn}{\varepsilon_0} \to E_r = \frac{qn}{2\varepsilon_0}r$$

Self-Magnetic Field by Ampere's Law

$$2\pi r B_{\theta} = \mu_0 q n \beta c \pi r^2 \rightarrow B_{\theta} = \mu_0 \frac{q n \beta c}{2} r$$





Brillouin Flow



Total Collective Force on beam particle

$$\vec{F} = q\left(\vec{E} + \vec{v} \times \vec{B}\right) \doteq \left(\frac{q^2 n}{2\varepsilon_0} \left(1 - \beta^2\right)\right) r$$

effective (de)focussing strength

$$k = \frac{\omega_p^2}{2\beta^2 c^2 \gamma^3} \qquad \text{where the non-relativistic "plasma frequency" is } \omega_p^2 = \frac{q^2 n}{\varepsilon_0 m}$$

By previous work with solenoids in the rotating frame, can have equilibrium (force balance) when

 $\frac{\omega_p^2}{2\gamma} = \Omega_c^2 \qquad \text{non-relativistic plasma and cyclotron frequencies } \omega_L = \frac{\Omega_c}{2\gamma}$

This state is known as **Brillouin Flow** and neglects

beam temperature (fluid flow)





Comments



• Some authors, Reiser in particular, define a relativistic plasma frequency

$$\omega_p^2 = \frac{q^2 n}{\varepsilon_0 \gamma^3 m}$$

• Lawson's book has a nice discussion about why it is impossible to establish a relativistic Brillouin flow in a device where beam is extracted from a single cathode at an equipotential surface. In this case one needs to have either sheering of the rotation or non-uniform density in the selfconsistent solution.







Vlasov-Poisson System

$$\frac{d\psi}{dt} = \frac{\partial\psi}{\partial t} + \dot{q}\frac{\partial\psi}{\partial x} + \dot{p}\frac{\partial\psi}{\partial p} = 0$$
$$\dot{q} = p / m$$
$$\dot{p}_i = q \left(E_i + \left(\vec{v} \times \vec{B}\right)_i \right)$$

• Self-consistent Field

$$\vec{\nabla} \cdot \vec{E} = -\nabla^2 \phi = \frac{en}{\varepsilon_0}$$
$$n = \int \psi d^3 \dot{\vec{q}}$$



Thomas Jefferson National Accelerator Facility



K-V Distribution



• Single value for the transverse Hamiltonian

$$\frac{1}{\varepsilon_x} \left(\frac{x^2 + (\alpha_x x + \beta_x x')^2}{\beta_x} \right) + \frac{1}{\varepsilon_y} \left(\frac{y^2 + (\alpha_y y + \beta_y y')^2}{\beta_y} \right) = C$$

$$\psi(x, x', y, y') \propto \delta(C - 1)$$

$$\rho(z) = \frac{I}{\pi \beta c X Y}$$

$$K = \frac{I}{I_0} \frac{2}{\beta^3 \gamma^3} \qquad I_0 = \frac{4\pi \varepsilon_0 m c^3}{q}$$

$$E_x = \frac{I}{\pi \varepsilon_0 \beta c} \frac{x}{Y(X + Y)}$$



Thomas Jefferson National Accelerator Facility Graduate Accelerator Physics Fall 2015



K-V Envelope Equation

 $E_{x} = \frac{I}{\pi \varepsilon_{0} \beta c} \frac{x}{X(X+Y)}$ $E_{y} = \frac{I}{\pi \varepsilon_{c} \beta c} \frac{y}{Y(X+Y)}$ $x'' + k_y x - \frac{2K}{X(X+Y)} x = 0$ $y'' + k_y x - \frac{2K}{Y(X+Y)} x = 0$ **Envelope Equation** $X'' + k_x X - \frac{2K}{(X+Y)} - \frac{\varepsilon_x^2}{X^3} = 0$ $Y'' + k_y Y - \frac{2K}{(X+Y)} - \frac{\varepsilon_y^2}{Y^3} = 0$



Thomas Jefferson National Accelerator Facility





Waterbag Distribution

• Lemons and Thode were first to point out SC field is solved as Bessel Functions for a certain equation of state. Later, others, including my advisor and I showed the equation of state was exact for the waterbag distribution.

$$H_{T} = \frac{p_{z}^{2}}{2m} (x'^{2} + y'^{2}) + \frac{m\omega_{0}^{2} (x^{2} + y^{2})}{2} + e\phi_{SC}$$

$$\psi = A\Theta (H_{0} - H_{T})$$

$$n(r) = \iint \psi dx' dy' = n_{0} \left(1 - \frac{m\omega_{0}^{2} (x^{2} + y^{2})}{2H_{0}} - \frac{\phi_{SC}}{\phi_{0}} \right) \quad \phi_{0} = H_{0} / e$$

$$\sigma_{v}^{2} = \frac{\iint \psi p_{z}^{2} (x'^{2} + y'^{2}) dx' dy'}{m^{2} \iint \psi dx' dy'} = \frac{H_{0}}{m} \left(1 - \frac{m\omega_{0}^{2} (x^{2} + y^{2})}{2H_{0}} - \frac{\phi_{SC}}{\phi_{0}} \right)$$



Thomas Jefferson National Accelerator Facility





Self-consistent potential solves

$$\nabla^2 \phi_{SC} + \frac{\phi_{SC}}{\lambda_D^2} = \frac{en_0}{\varepsilon_0} \left[\frac{m\omega_0^2 \left(x^2 + y^2 \right)}{2H_0} - 1 \right]$$
$$\lambda_D = \frac{\sigma_v}{\omega_p} = \sqrt{\frac{\varepsilon_0 m}{e^2 n_0} \frac{H_0}{m}} = \sqrt{\frac{\varepsilon_0 H_0}{e^2 n_0}} \quad \text{Debye Length}$$

Analytic solutions in terms of Modified Bessel Functions

$$e\phi(r) = \frac{m\omega_0^2(x^2 + y^2)}{2} + A(I_0(r/\lambda_D) - 1) + BK_0(r/\lambda_D)$$

B = 0 by boundary condition

A chosen so that solution without I_0 solution to inhomogeneous eqn.





Equation for Beam Radius



$$n_b(r) = \hat{n}_b \frac{I_0(r_b / \lambda_D) - I_0(r / \lambda_D)}{I_0(r_b / \lambda_D) - 1}$$



Thomas Jefferson National Accelerator Facility



Luminosity and Beam-Beam Effect

- Luminosity Defined
- Beam-Beam Tune Shift
- Luminosity Tune-shift Relationship (Krafft-Ziemann Thm.)
- Beam-Beam Effect





Events per Beam Crossing

 In a nuclear physics experiment with a beam crossing through a thin fixed target



- Probability of single event, per beam particle passage is $P = n\sigma l$
- σ is the "cross section" for the process (area units)







• Probability an event is generated by a single particle of Beam 1 crossing Beam 2 bunch with Gaussian density*

$$P = \sigma \frac{N_{2} \exp(-x^{2} / 2\sigma_{2x}^{2}) \exp(-y^{2} / 2\sigma_{2y}^{2})}{(2\pi)^{3/2} \sigma_{2x} \sigma_{2y} \sigma_{2z}} \int_{-\infty}^{\infty} \exp(-z^{2} / 2\sigma_{2z}^{2}) dz$$
$$= \frac{N_{2} \exp(-x^{2} / 2\sigma_{2x}^{2}) \exp(-y^{2} / 2\sigma_{2y}^{2})}{2\pi \sigma_{2x} \sigma_{2y}} \sigma$$

* This expression still correct when relativity done properly



Thomas Jefferson National Accelerator Facility



Collider Luminosity

Probability an event is generated by a Beam 1 bunch with Gaussian density crossing a Beam 2 bunch with Gaussian density

$$P = \frac{N_1 N_2}{2\pi \sqrt{\sigma_{1x}^2 + \sigma_{2x}^2} \sqrt{\sigma_{1y}^2 + \sigma_{2y}^2}} \sigma$$

• Event rate with equal transverse beam sizes

$$\frac{dN}{dt} = \frac{fN_1N_2}{4\pi\sigma_x\sigma_y}\sigma = \mathcal{L}\sigma$$

• Luminosity

$$\mathcal{L} = \frac{fN_1N_2}{4\pi\sigma_x\sigma_y} \sim 10^{33} \text{sec}^{-1} \text{cm}^{-2},$$

for f = 100 MHz, $N_1 = N_2 = 10^{10}$, $\sigma_x = \sigma_y = 10$ microns



Thomas Jefferson National Accelerator Facility



Beam-Beam Tune Shift



- As we've seen previously, in a ring accelerator the number of transverse oscillations a particle makes in one circuit is called the "betatron tune" *Q*.
- Any deviation from the design values of the tune (in either the horizontal or vertical directions), is called a "tune shift". For long term stability of the beam in a ring accelerator, the tune must be highly controlled.

$$M_{tot} = \begin{pmatrix} 1 & 0 \\ -1/f & 1 \end{pmatrix} \begin{pmatrix} \cos \mu & \beta^* \sin \mu \\ -\sin \mu / \beta^* & \cos \mu \end{pmatrix}$$
$$= \begin{pmatrix} \cos \mu & \beta^* \sin \mu \\ -\cos \mu / f - \sin \mu / \beta^* & \cos \mu - (\beta^* / f) \sin \mu \end{pmatrix}$$



Thomas Jefferson National Accelerator Facility





$$\cos\left(\mu + \Delta\mu\right) = \frac{\operatorname{Tr}\left(M_{tot}\right)}{2} = \cos\mu - \frac{\beta^{*}}{2f}\sin\mu$$
$$\xi = \Delta Q = \frac{\Delta\mu}{2\pi} = \frac{\beta^{*}}{4\pi f} \qquad \beta^{*} \ll f$$



Thomas Jefferson National Accelerator Facility



Bessetti-Erskine Solution

• 2-D potential of Bi-Gaussian transverse distribution

$$\rho(x, y) = \frac{Q'}{2\pi\sigma_x \sigma_y} \exp\left(-\frac{x^2}{2\sigma_x^2}\right) \exp\left(-\frac{y^2}{2\sigma_y^2}\right)$$

• Potential Theory gives solution to Poisson Equation

$$\nabla^2 \phi = \frac{\rho(x, y)}{\varepsilon_0}$$
$$(x, y) = \frac{Q'}{4\pi\varepsilon_0} \int_0^\infty \frac{\exp\left(-\frac{x^2}{2\sigma_x^2 + q}\right) \exp\left(-\frac{y^2}{2\sigma_y^2 + q}\right)}{\sqrt{2\sigma_x^2 + q}\sqrt{2\sigma_y^2 + q}} dq$$

• Bassetti and Erskine manipulate this to



 ϕ



$$E_{x} = \frac{Q'}{2\varepsilon_{0}\sqrt{2\pi(\sigma_{x}^{2} - \sigma_{y}^{2})}} \operatorname{Im}\left[w\left(\frac{x + iy}{\sqrt{2(\sigma_{x}^{2} - \sigma_{y}^{2})}}\right) - \exp\left(-\frac{x^{2}}{2\sigma_{x}^{2}} - \frac{x^{2}}{2\sigma_{y}^{2}}\right)w\left(\frac{x\left(\frac{\sigma_{y}}{\sigma_{x}}\right) + iy\left(\frac{\sigma_{x}}{\sigma_{y}}\right)}{\sqrt{2(\sigma_{x}^{2} - \sigma_{y}^{2})}}\right)\right]$$

$$E_{y} = \frac{Q'}{2\varepsilon_{0}\sqrt{2\pi(\sigma_{x}^{2} - \sigma_{y}^{2})}} \operatorname{Re}\left[w\left(\frac{x + iy}{\sqrt{2(\sigma_{x}^{2} - \sigma_{y}^{2})}}\right) - \exp\left(-\frac{x^{2}}{2\sigma_{x}^{2}} - \frac{x^{2}}{2\sigma_{y}^{2}}\right)w\left(\frac{x\left(\frac{\sigma_{y}}{\sigma_{x}}\right) + iy\left(\frac{\sigma_{x}}{\sigma_{y}}\right)}{\sqrt{2(\sigma_{x}^{2} - \sigma_{y}^{2})}}\right)\right]$$

$$w(z) \quad \text{Complex error function}$$

• We need 2-D linear field for small displacements

$$E_{x}(x,0) = -\frac{\partial \phi}{\partial x} \doteq \frac{Q'x}{2\pi\varepsilon_{0}} \int_{0}^{\infty} \frac{1}{\left(\sqrt{2\sigma_{x}^{2}+q}\right)^{3/2} \sqrt{2\sigma_{y}^{2}+q}} dq$$



Thomas Jefferson National Accelerator Facility



-



• Can do the integral analytically



• Similarly for the *y*-direction

$$E_{y}(0, y) = -\frac{\partial \phi}{\partial y} \doteq \frac{Q'y}{2\pi\varepsilon_{0}} \frac{1}{\sigma_{y}(\sigma_{x} + \sigma_{y})}$$





Linear Beam-Beam Kick

• Linear kick received after interaction with bunch

$$\Delta(\gamma_1\beta_{1x}mc) = q_1 \int_{-\infty}^{\infty} \left(\vec{E}_{2x} + \left(\vec{v} \times \vec{B}\right)_{2x}\right) \left(\vec{x}_1(t), t\right) dt$$

by relativity, for oppositely moving beams

$$\Delta \gamma \beta_{1x} mc = q_1 \left(1 + \beta_{1z} \beta_{2z} \right) \int_{-\infty}^{\infty} \left(\vec{E}_{2x} \right) \left(\vec{x}_1(t), t \right) dt$$

Following linear Bassetti-Erskine model

$$E_{2x}(x,0,z,t) = \frac{q_2 x}{2\pi\varepsilon_0} \frac{1}{\sigma_x(\sigma_x + \sigma_y)} \frac{1}{\sqrt{2\pi}\sigma_2} \exp\left(-\frac{\left(z - \beta_{2z}ct\right)^2}{2\sigma_z^2}\right)$$

$$q_1 \text{ moves with } \vec{x}(t) = (x,0,-\beta_{1z}ct)$$







Linear Beam-Beam Tune Shift

$$\therefore \Delta \gamma \beta_{1x} mc = q_1 \frac{(1 + \beta_{1z} \beta_{2z})}{\beta_{1z} + \beta_{2z}} \frac{q_2 x}{2\pi \varepsilon_0 c} \frac{1}{\sigma_x (\sigma_x + \sigma_y)}$$

$$1/f = \frac{2N_2}{\gamma_1} \frac{(1 + \beta_{1z} \beta_{2z})}{\beta_{1z} + \beta_{2z}} \frac{r_1}{\sigma_x (\sigma_x + \sigma_y)} \qquad r_1 = \frac{e^2}{4\pi \varepsilon_0 m_1 c^2}$$

$$1/f \doteq \frac{2N_2 r_1}{\gamma_1 \sigma_x (\sigma_x + \sigma_y)} \qquad \text{Both beams relativistic}$$

From linear Bassetti-Erskine model, and replacing the beam size

$$\xi_x^1 = \frac{N_2 r_1}{2\pi \gamma_1} \frac{1}{\varepsilon_x^1 \left(1 + \sigma_y / \sigma_x\right)} \qquad \xi_y^1 = \frac{N_2 r_1}{2\pi \gamma_1} \frac{1}{\varepsilon_y^i \left(1 + \sigma_y / \sigma_x\right) \left(\sigma_x / \sigma_y\right)}$$

Argument entirely symmetric wrt choice of bunch 1 and 2

$$\xi_x^i = \frac{N_{\bar{i}}r_i}{2\pi\gamma_i} \frac{1}{\varepsilon_x^i \left(1 + \sigma_y / \sigma_x\right)} \qquad \xi_y^i = \frac{N_{\bar{i}}r_i}{2\pi\gamma_i} \frac{1}{\varepsilon_y^i \left(1 + \sigma_y / \sigma_x\right) \left(\sigma_x / \sigma_y\right)}$$

Jefferson Lab

Thomas Jefferson National Accelerator Facility Graduate Accelerator Physics Fall 2015



Luminosity Beam-Beam tune-shift () P

• Express Luminosity in terms of the (larger!) vertical tune shift (*i* either 1 or 2)

$$\mathcal{L} = \frac{f N_i \xi_y^i \gamma_i}{2r_i \beta_{iy}^*} \left(1 + \sigma_y / \sigma_x\right) = \frac{I_i}{e} \frac{\xi_y^i \gamma_i}{2r_i \beta_{iy}^*} \left(1 + \sigma_y / \sigma_x\right)$$

- Necessary, but not sufficient, for self-consistent design
- Expressed in this way, and given a known limit to the beam-beam tune shift, the only variables to manipulate to increase luminosity are the stored current, the aspect ratio, and the β^* (beta function value at the interaction point)
- Applies to ERL-ring colliders, stored beam (ions) only





Luminosity-Deflection Theorem

- Luminosity-tune shift formula is linearized version of a much more general formula discovered by Krafft and generalized by V. Ziemann.
- Relates easy calculation (luminosity) to a hard calculation (beam-beam force), and contains all the standard results in beam-beam interaction theory.
- Based on the fact that the relativistic beam-beam force is almost entirely transverse, i. e., 2-D electrostatics applies.





2-D Electrostatics Theorem

$$\vec{E}(\vec{x}) = \frac{2Q'}{4\pi\varepsilon_0} \frac{\vec{x} - \vec{x}'}{|\vec{x} - \vec{x}'|^2}$$

$$\vec{F}_{21}' = -\vec{F}_{12}' = \frac{1}{2\pi\varepsilon_0} \iint \rho_2(\vec{x}_2) \frac{\vec{x}_2 - \vec{x}_1}{|\vec{x}_2 - \vec{x}_1|} \rho_1(\vec{x}_1) d^2 \vec{x}_1 d^2 \vec{x}_2 \quad 1 \text{ on } 2$$

$$n_1(\vec{x}_1) = \rho_1(\vec{x}_1) / Q_1' \quad n_2(\vec{x}_2) = \rho_1(\vec{x}_2 + \vec{b}) / Q_1' \text{ zero centerred}$$

$$Q_i' = \iint \rho_i(\vec{x}) d^2 \vec{x} \qquad \vec{b} = \iint \vec{x} \rho_2(\vec{x}) d^2 \vec{x} / Q_2'$$

$$\vec{F}_{21}' = -\vec{F}_{12}' = \frac{Q_1' Q_2'}{2\pi\varepsilon_0} \iint n_2(\vec{x}_2) \frac{\vec{x}_1 + \vec{b} - \vec{x}_2}{|\vec{x}_1 + \vec{b} - \vec{x}_2|^2} n_1(\vec{x}_1) d^2 \vec{x}_1 d^2 \vec{x}_2$$



Thomas Jefferson National Accelerator Facility



)))



$$\vec{\nabla}_{\vec{b}} \cdot \frac{\vec{x}_1 + \vec{b} - \vec{x}_2}{\left|\vec{x}_1 + \vec{b} - \vec{x}_2\right|^2} = 2\pi\delta\left(x_2 + b_x + x_1\right)\delta\left(y_2 + b_y + y_1\right)$$
$$\vec{\nabla}_{\vec{b}} \cdot \frac{\vec{F}}{\left|\vec{x}_1 + \vec{b} - \vec{x}_2\right|^2} = (\vec{x} + \vec{b})e_x(\vec{x})d^2\vec{x}$$

$$\vec{\nabla}_{\vec{b}} \cdot \vec{F}_{21}' = \frac{1}{\varepsilon_0} \iint \rho_2 \left(\vec{x} + \vec{b} \right) \rho_1 \left(\vec{x} \right) d^2 \vec{x}$$

Generalizes
$$\vec{\nabla} \cdot \vec{E} = \frac{\rho}{\varepsilon_0} \quad \left(\text{take } \rho_2 \left(\vec{x} \right) \propto \delta^2 \left(\vec{x} + \vec{b} \right) \right)$$

Transverse interaction in the beam-beam problem

$$\Delta p_1 = \frac{q_1 q_2}{2\pi\varepsilon_0 c} \frac{\vec{x}_1 - \vec{x}_2}{\left|\vec{x}_1 - \vec{x}_2\right|^2}$$



Thomas Jefferson National Accelerator Facility





$$\begin{aligned} \vec{D}(\vec{b}) &= \Delta \gamma_1 \vec{\beta}_1 = -\Delta m_2 \gamma_2 \vec{\beta}_2 / m_1 \\ &= \frac{q_1 q_2}{m_1 c^2} \iint n_2 (\vec{x}_2) \frac{\vec{x}_1 - \vec{x}_2 - \vec{b}}{\left| \vec{x}_1 - \vec{x}_2 - \vec{b} \right|^2} n_1 (\vec{x}_1) d^2 \vec{x}_1 d^2 \vec{x}_2 \\ \vec{\nabla}_{\vec{b}} \cdot \vec{D}(\vec{b}) &= 4\pi N_2 r_e \iint n_2 (\vec{x} - \vec{b}) n_1 (\vec{x}) d^2 \vec{x} \quad r_e = \frac{e^2}{4\pi \varepsilon_0 m c^2} \\ L(\vec{b}) &= N_1 N_2 \iint n_2 (\vec{x} - \vec{b}) n_1 (\vec{x}) d^2 \vec{x} \\ L(\vec{b}) &= \frac{N_1}{4\pi r_e} \vec{\nabla}_{\vec{b}} \cdot \vec{D}(\vec{b}) \end{aligned}$$

$$L(\vec{b}) = -\frac{N_2}{4\pi r_e} \vec{\nabla}_{\vec{b}} \cdot \left(\Delta \gamma_2 \vec{\beta}_2\right)$$



Thomas Jefferson National Accelerator Facility





$$\begin{pmatrix} D_x \\ D_y \end{pmatrix} = \frac{\gamma_1}{2f} \begin{pmatrix} \sigma_y / \sigma_x & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} b_x \\ b_y \end{pmatrix}$$

$$N_1 \gamma \xi (1 - \gamma) = 1 - \xi$$

$$L = \frac{N_1 \gamma \varsigma}{2r_e \beta^*} \left(1 + \sigma_y / \sigma_x \right) \quad \text{as before}$$

Maximum when

 $\frac{\partial}{\partial b_x} \left[\frac{\partial D_x}{\partial b_x} \right] = 0, \qquad \frac{\partial}{\partial b_y} \left[\frac{\partial D}{\partial b_y} \right] = 0$



Thomas Jefferson National Accelerator Facility





Luminosity-Deflection Pairs

• Round Beam Fast Model

$$\vec{D}(\vec{b}) = \frac{2N_2 r_e \vec{b}}{\sigma^2 + b^2} \qquad L(\vec{b}) = \frac{N_1 N_2 \sigma^2}{\pi (\sigma^2 + b^2)^2}$$
• Gaussian Macroparticles

$$\vec{D}(\vec{b}) = \vec{D}_{Bassetti_Erskine} \left(\vec{b}; \sqrt{\sigma_{1x}^2 + \sigma_{2x}^2}; \sqrt{\sigma_{1y}^2 + \sigma_{2y}^2}\right)$$

$$L(\vec{b}) = \frac{N_1 N_2}{2\pi \sqrt{\sigma_{1x}^2 + \sigma_{2x}^2} \sqrt{\sigma_{1y}^2 + \sigma_{2y}^2}} \exp\left(-\frac{b_x^2}{\sqrt{\sigma_{1x}^2 + \sigma_{2x}^2}}\right) \exp\left(-\frac{b_y^2}{\sqrt{\sigma_{1y}^2 + \sigma_{2y}^2}}\right)$$
• Smith-Laslett Model

$$\vec{D}(\vec{b}) = \frac{2N_2 r_e \vec{b}}{\sqrt{(4b^2 + 2b^4)}} - \frac{4b^2}{2\pi \sqrt{(4b^2 + 2b^4)}} \left\{\sinh^{-1}\left[\frac{b^3}{2} + \frac{3b}{2}\right] + \sinh^{-1}\left[\frac{b}{2}\right]\right\}$$

$$\begin{split} \vec{D}(\vec{b}) &= \frac{2N_2 T_e \vec{b}}{\hat{b}^2 A B} \left\{ \frac{(1-1)^2}{(4\hat{b}^2 + \hat{b}^4)^2} - \frac{4\vec{b}}{(4\hat{b}^2 + \hat{b}^4)^{3/2}} \left\{ \sinh^{-1} \left[\frac{\vec{b}}{2} + \frac{3\vec{b}}{2} \right] + \sinh^{-1} \left[\frac{\vec{b}}{2} \right] \right\} \right\} \\ \vec{L}(\vec{b}) &= \frac{N_1 N_2}{\pi A B} \left\{ \frac{(2\hat{b}^2 - 4)\hat{b}^2}{(4\hat{b}^2 + \hat{b}^4)^2} - \frac{4\hat{b}^2 (1 + \hat{b}^2)}{(4\hat{b}^2 + \hat{b}^4)^{5/2}} \left\{ \sinh^{-1} \left[\frac{\hat{b}^3}{2} + \frac{3\hat{b}}{2} \right] + \sinh^{-1} \left[\frac{\hat{b}}{2} \right] \right\} \right\} \end{split} \\ \vec{b}^2 &= \left(\frac{b_x}{A} \right)^2 + \left(\frac{b_y}{B} \right)^2 \end{split}$$



Thomas Jefferson National Accelerator Facility

