



Accelerator Physics

Synchrotron Radiation

A. Bogacz, G. A. Krafft, and T. Zolkin

Jefferson Lab

Colorado State University

Lecture 8

Synchrotron Radiation



Accelerated particles emit electromagnetic radiation. Emission from very high energy particles has unique properties for a radiation source. As such radiation was first observed at one of the earliest electron synchrotrons, radiation from high energy particles (mainly electrons) is known generically as synchrotron radiation by the accelerator and HENP communities.

The radiation is highly collimated in the beam direction

From relativity

$$ct' = \gamma ct - \gamma\beta z$$

$$x' = x$$

$$y' = y$$

$$z' = -\gamma\beta ct + \gamma z$$



Lorentz invariance of wave phase implies $k^\mu = (\omega/c, k_x, k_y, k_z)$ is a Lorentz 4-vector

$$\omega' = \gamma\omega - \gamma\beta k_z c$$

$$k'_x = k_x$$

$$k'_y = k_y$$

$$k'_z = -\gamma\beta\omega / c + \gamma k_z$$

$$\sin \theta = \frac{\sqrt{k_x^2 + k_y^2}}{\omega / c}$$

$$\sin \theta' = \frac{\sqrt{k_x'^2 + k_y'^2}}{\omega' / c}$$

$$\cos \theta' = \frac{k'_z}{\omega' / c}$$

$$\omega / c = \gamma\omega' / c + \gamma\beta k'_z = \gamma(1 + \beta \cos \theta')(\omega' / c)$$

$$\omega' / c = \gamma\omega / c - \gamma\beta k_z = \gamma(1 - \beta \cos \theta)(\omega / c)$$



$$\theta \approx \sin \theta = \frac{\sin \theta'}{\gamma (1 + \beta \cos \theta')}$$

Therefore all radiation with $\theta' < \pi / 2$, which is roughly $1/2$ of the photon emission for dipole emission from a transverse acceleration in the beam frame, is Lorentz transformed into an angle less than $1/\gamma$. Because of the strong Doppler shift of the photon energy, higher for $\theta \rightarrow 0$, most of the energy in the photons is within a cone of angular extent $1/\gamma$ around the beam direction.

Larmor's Formula



For a particle executing non-relativistic motion, the total power emitted in electromagnetic radiation is (Larmor, verified later)

$$P(t) = \frac{1}{6\pi\epsilon_0} \frac{q^2}{c^3} |\vec{a}|^2 = \frac{1}{6\pi\epsilon_0} \frac{e^2}{m^2 c^3} |\dot{\vec{p}}|^2$$

Lienard's relativistic generalization: Note both dE and dt are the fourth component of relativistic 4-vectors when one is dealing with photon emission. Therefore, their ratio must be a Lorentz invariant. The invariant that reduces to Larmor's formula in the non-relativistic limit is

$$P = - \frac{e^2}{6\pi\epsilon_0 c} \frac{du^\mu}{d\tau} \frac{du_\mu}{d\tau}$$



$$P(t) = \frac{e^2}{6\pi\epsilon_0 c} \gamma^6 \left(\dot{\vec{\beta}}^2 - \left[\vec{\beta} \times \dot{\vec{\beta}} \right]^2 \right)$$

For acceleration along a line, second term is zero and first term for the radiation reaction is small compared to the acceleration as long as gradient less than 10^{14} MV/m. Technically impossible.

For transverse bend acceleration $\dot{\vec{\beta}} = -\frac{\beta^2 c}{\rho} \hat{r}$

$$P(t) = \frac{e^2 c}{6\pi\epsilon_0 \rho^2} \beta^4 \gamma^4$$

Fractional Energy Loss



$$\delta E = \frac{e^2}{6\pi\epsilon_0\rho} \Theta \beta^3 \gamma^4$$

For one turn with isomagnetic bending fields

$$\frac{\delta E}{E_{beam}} = \frac{4\pi r_e}{3\rho} \beta^3 \gamma^3$$

r_e is the classical electron radius: $2.82 \cdot 10^{-13}$ cm

Radiation Power Distribution



Consulting your favorite Classical E&M text (Jackson, Schwinger, Landau and Lifshitz Classical Theory of Fields)

$$\frac{dP}{d\omega} = \frac{\sqrt{3}}{8\pi^2 \epsilon_0} \frac{e^2}{\rho} \gamma \frac{\omega}{\omega_c} \int_{\omega/\omega_c}^{\infty} K_{5/3}(x) dx$$

Critical Frequency




Critical (angular) frequency is

$$\omega_c = \frac{3}{2} \gamma^3 \frac{c}{\rho}$$

Energy scaling of critical frequency is understood from $1/\gamma$ emission cone and fact that $1 - \beta \sim 1/(2\gamma^2)$

$$t_A = -\frac{\rho}{\gamma\beta c}$$

$$t_B = \frac{\rho}{\gamma\beta c} \approx \frac{\rho}{\gamma c} + \frac{\rho}{2\gamma^3 c}$$



Photon Number



$$P = \int_0^{\infty} \frac{dP}{d\omega} d\omega = \frac{\sqrt{3}}{8\pi^2 \epsilon_0} \frac{e^2}{\rho} \omega_c \gamma \int_0^{\infty} \xi \int_{\xi}^{\infty} K_{5/3}(x) dx d\xi = \frac{e^2 c}{6\pi \epsilon_0 \rho^2} \gamma^4$$

$$\frac{d\dot{n}}{d\omega} = \frac{1}{\hbar \omega} \frac{dP}{d\omega}$$

$$\langle \hbar \omega \rangle = \frac{\int_0^{\infty} \hbar \omega \frac{d\dot{n}}{d\omega} d\omega}{\int_0^{\infty} \frac{d\dot{n}}{d\omega} d\omega} = \frac{8}{15\sqrt{3}} \hbar \omega_c$$

$$\dot{n} = \frac{5\alpha}{2\sqrt{3}} \frac{c}{\rho} \gamma \quad \delta n = \frac{5\alpha}{2\sqrt{3}} \Theta \gamma \quad \alpha = \frac{e^2}{4\pi \epsilon_0 \hbar c} \approx \frac{1}{137}$$

Insertion Devices (ID)



Often periodic magnetic field magnets are placed in beam path of high energy storage rings. The radiation generated by electrons passing through such insertion devices has unique properties.

Field of the insertion device magnet

$$\vec{B}(x, y, z) = B(z) \hat{y} \quad B(z) \approx B_0 \cos(2\pi z / \lambda_{ID})$$

Vector potential for magnet (1 dimensional approximation)

$$\vec{A}(x, y, z) = A(z) \hat{x} \quad A(z) \approx \frac{B_0 \lambda_{ID}}{2\pi} \sin(2\pi z / \lambda_{ID})$$

Electron Orbit



Uniformity in x -direction means that canonical momentum in the x -direction is conserved

$$v_x(z) = \frac{eA(z)}{\gamma m} = \frac{K}{\gamma} c \sin(2\pi z / \lambda_{ID})$$

$$x(z) = \int \frac{v_x}{v_z} dz \approx -\frac{1}{\langle \beta_z \rangle} \frac{K}{\gamma} \frac{\lambda_{ID}}{2\pi} \cos(2\pi z / \lambda_{ID})$$

Field Strength Parameter

$$K \equiv \frac{eB_0 \lambda_{ID}}{2\pi m c}$$

Average Velocity



Energy conservation gives that γ is a constant of the motion

$$\beta_z(z) = \sqrt{1 - \frac{1}{\gamma^2} - \beta_x^2(z)} = \sqrt{\beta_{z0}^2 - \beta_x^2(z)}$$

Average longitudinal velocity in the insertion device is

$$\beta^{*2} = \langle \beta_z \rangle^2 = 1 - \frac{1}{\gamma^2} - \frac{K^2}{2\gamma^2}$$

Average rest frame has

$$\gamma^{*2} = \frac{1}{1 - \beta^{*2}} = \frac{\gamma^2}{1 + K^2/2}$$

Relativistic Kinematics



In average rest frame the insertion device is Lorentz contracted, and so its wavelength is

$$\lambda^* = \lambda_{ID} / \beta^* \gamma^*$$

The sinusoidal wiggling motion emits with angular frequency

$$\omega^* = 2\pi c / \lambda^*$$

Lorentz transformation formulas for the wave vector of the emitted radiation

$$k^* = \gamma^* k (1 - \beta^* \cos \theta)$$

$$k_x^* = k_x = k \sin \theta \cos \varphi$$

$$k_y^* = k_y = k \sin \theta \sin \varphi$$

$$k_z^* = \gamma^* k (\cos \theta - \beta^*)$$

ID (or FEL) Resonance Condition



Angle transforms as

$$\cos \theta^* = \frac{k_z^*}{k^*} = \frac{(\cos \theta - \beta^*)}{(1 - \beta^* \cos \theta)}$$

Wave vector in lab frame has

$$k = \frac{k^*}{\gamma^* (1 - \beta^* \cos \theta)} = \frac{2\pi\beta^* c}{\lambda_{ID} (1 - \beta^* \cos \theta)}$$

In the forward direction $\cos \theta = 1$

$$\lambda_e \approx \frac{\lambda_{ID}}{2\gamma^{*2}} = \frac{\lambda_{ID}}{2\gamma^2} (1 + K^2 / 2)$$

Power Emitted Lab Frame



Larmor/Lienard calculation in the lab frame yields

$$\langle P \rangle = \frac{e^2}{6\pi\epsilon_0} \gamma^4 \beta_{z0}^2 c \left(\frac{K}{\gamma} \right)^2 \left(\frac{2\pi}{\lambda_{ID}} \right)^2 \frac{1}{2}$$

Total energy radiated after one passage of the insertion device

$$\delta E = 2\pi^2 \frac{e^2}{6\pi\epsilon_0 \lambda_{ID}} \gamma^2 \frac{\beta_{z0}^2}{\beta^*} NK^2$$

Power Emitted Beam Frame



Larmor/Lienard calculation in the beam frame yields

$$\langle P^* \rangle = \frac{e^2}{6\pi\epsilon_0} cK^2 \left(\frac{2\pi}{\lambda^*} \right)^2 \frac{1}{2}$$

Total energy of each photon is $\hbar 2\pi c/\lambda^*$, therefore the total number of photons radiated after one passage of the insertion device

$$N_\gamma = \frac{2\pi}{3} \alpha N K^2$$

Spectral Distribution in Beam Frame



Begin with power distribution in beam frame: dipole radiation pattern (single harmonic only when $K \ll 1$; replace γ^* by γ , β^* by β)

$$\frac{dP^*}{d\Omega^*} = \frac{e^2 c}{32\pi^2 \epsilon_0} K^2 k^{*2} \sin^2 \Theta^*$$

Number distribution in terms of wave number

$$\frac{dN_\gamma}{d\Omega^*} = \frac{\alpha}{4} NK^2 \frac{k_y^{*2} + k_z^{*2}}{k^{*2}}$$

Solid angle transformation

$$d\Omega^* = \frac{d\Omega}{\gamma^2 (1 - \beta \cos \theta)^2}$$



Number distribution in beam frame

$$\frac{dN_\gamma}{d\Omega} = \frac{\alpha}{4} NK^2 \frac{\sin^2 \theta \sin^2 \varphi + \gamma^2 (\cos \theta - \beta)^2}{\gamma^4 (1 - \beta \cos \theta)^4}$$

Energy is simply

$$E(\theta) = \hbar \frac{2\pi\beta c}{\lambda_{ID} (1 - \beta \cos \theta)} \quad \hat{E}(\theta) = \frac{1}{(1 - \beta \cos \theta)}$$

Number distribution as a function of normalized lab-frame energy

$$\frac{dN_\gamma}{d\hat{E}} = \frac{\alpha\pi}{4\gamma^2\beta^3} NK^2 \left[\left(\frac{\hat{E}}{\gamma^2} - 1 \right)^2 + \beta^2 \right]$$

Average Energy




Limits of integration

$$\cos \theta = 1 \quad \hat{E} = \frac{1}{1 - \beta} \quad \cos \theta = -1 \quad \hat{E} = \frac{1}{1 + \beta}$$

Average energy is also analytically calculable

$$\langle E \rangle = \frac{\int_0^{\infty} E \frac{dN_{\gamma}}{d\hat{E}} d\hat{E}}{\int_0^{\infty} \frac{dN_{\gamma}}{d\hat{E}} d\hat{E}} = \gamma^2 \hbar 2\pi\beta c / \lambda_{ID} \approx \frac{E_{\max}}{2}$$

Number Spectrum



Conventions on Fourier Transforms



For the time dimensions

$$\tilde{f}(\omega) = \int_{-\infty}^{\infty} f(t) e^{i\omega t} dt$$

$$f(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \tilde{f}(\omega) e^{-i\omega t} d\omega$$

Results on Dirac delta functions

$$\tilde{\delta}(\omega) = \int_{-\infty}^{\infty} \delta(t) e^{i\omega t} dt = 1$$

$$\delta(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-i\omega t} d\omega$$



For the three spatial dimensions

$$\tilde{f}(\vec{k}) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(\vec{x}) e^{-i\vec{k} \cdot \vec{x}} d^3 \vec{x}$$

$$f(\vec{x}) = \frac{1}{(2\pi)^3} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \tilde{f}(\vec{k}) e^{+i\vec{k} \cdot \vec{x}} d^3 \vec{k}$$

$$\delta^3(\vec{x}) = \delta(\vec{x}) = \frac{1}{(2\pi)^3} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{+i\vec{k} \cdot \vec{x}} d^3 \vec{k}$$

Green Function for Wave Equation



Solution to inhomogeneous wave equation

$$\left[\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \right] G(\vec{x}, t; \vec{x}', t')$$
$$= -4\pi \delta(\vec{x} - \vec{x}') \delta(t - t')$$

Will pick out the solution with causal boundary conditions, i. e.

$$G(\vec{x}, t; \vec{x}', t') = 0 \quad t < t'$$

This choice leads automatically to the so-called *Retarded* Green Function



In general

$$G(\vec{x}, t; \vec{x}', t') = 0 \quad t < t'$$

$$G(\vec{x}, t; \vec{x}', t') = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \left[A(\vec{k}) e^{i(\vec{k} \cdot \vec{x} - \omega t)} + B(\vec{k}) e^{i(\vec{k} \cdot \vec{x} + \omega t)} \right] d^3 \vec{k} \quad t > t'$$

because there are two possible signs of the frequency for each value of the wave vector. To solve the homogeneous wave equation it is necessary that

$$\omega(\vec{k}) = |\vec{k}|c$$

i.e., there is no dispersion in free space.



Continuity of G implies

$$A(\vec{k})e^{-i\omega t'} = -B(\vec{k})e^{i\omega t'}$$

Integrate the inhomogeneous equation between $t = t' + \varepsilon$ and $t = t' - \varepsilon$

$$\begin{aligned} -\frac{1}{c^2} \frac{\partial G(\vec{x}, t; \vec{x}', t')}{\partial t} \Big|_{t'+\varepsilon} &= -4\pi\delta(\vec{x} - \vec{x}') \\ \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \left[-i\omega A(\vec{k})e^{i(\vec{k}\cdot\vec{x} - \omega t')} + i\omega B(\vec{k})e^{i(\vec{k}\cdot\vec{x} + \omega t')} \right] d^3\vec{k} & \\ &= 4\pi c^2 \delta(\vec{x} - \vec{x}') \end{aligned}$$

$$A(\vec{k}) = -\frac{c^2}{(2\pi)^2 i\omega} e^{-i\vec{k}\cdot\vec{x}'} e^{i\omega t'}$$



$$\begin{aligned} G(\vec{x}, t; \vec{x}', t') &= \\ &= -\frac{c^2}{(2\pi)^2} \frac{1}{i} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{1}{\omega} \left[e^{i(\vec{k} \cdot (\vec{x} - \vec{x}') - \omega(t - t'))} - e^{i(\vec{k} \cdot (\vec{x} - \vec{x}') + \omega(t - t'))} \right] d^3 \vec{k} \\ & \qquad \qquad \qquad t > t' \\ &= \frac{c}{2\pi} \int_{-\infty}^{\infty} \frac{e^{ik|\vec{x} - \vec{x}'|}}{|\vec{x} - \vec{x}'|} e^{-i\omega(t - t')} dk - \frac{c}{2\pi} \int_{-\infty}^{\infty} \frac{e^{ik|\vec{x} - \vec{x}'|}}{|\vec{x} - \vec{x}'|} e^{+i\omega(t - t')} dk \quad t > t' \\ &= \frac{\delta(|\vec{x} - \vec{x}'| / c - t + t')}{|\vec{x} - \vec{x}'|} + 0 \end{aligned}$$

Called retarded because the influence at time t is due to the source evaluated at the retarded time

$$t' = t - |\vec{x} - \vec{x}'| / c$$

Retarded Solutions for Fields



$$\left[\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \right] \phi = -\frac{\rho}{\epsilon_0}$$

$$\left[\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \right] \vec{A} = -\mu_0 \vec{J}$$

$$\phi(\vec{x}, t) = \frac{1}{4\pi\epsilon_0} \int d^3x' dt' \frac{\rho(\vec{x}', t')}{|\vec{x} - \vec{x}'|} \delta(|\vec{x} - \vec{x}'|/c - t + t')$$

$$\vec{A}(\vec{x}, t) = \frac{\mu_0}{4\pi} \int d^3x' dt' \frac{\vec{J}(\vec{x}', t')}{|\vec{x} - \vec{x}'|} \delta(|\vec{x} - \vec{x}'|/c - t + t')$$

Tip: Leave the delta function in its integral form to do derivations.
Don't have to remember complicated delta-function rules

Delta Function Representation



$$\phi(\vec{x}, t) = \frac{1}{8\pi^2 \epsilon_0} \int d^3x' dt' d\omega \frac{\rho(\vec{x}', t')}{|\vec{x} - \vec{x}'|} e^{i\omega[|\vec{x} - \vec{x}'|/c - (t - t')]}$$

$$\vec{A}(\vec{x}, t) = \frac{\mu_0}{8\pi^2} \int d^3x' dt' d\omega \frac{\vec{J}(\vec{x}', t')}{|\vec{x} - \vec{x}'|} e^{i\omega[|\vec{x} - \vec{x}'|/c - (t - t')]}$$

Evaluation can be expedited by noting and using the symmetry of the Green function and using relations such as

$$\frac{\partial}{\partial t} f(t - t') = -\frac{\partial}{\partial t'} f(t - t')$$
$$\frac{\partial}{\partial \vec{x}} f(|\vec{x} - \vec{x}'|) = -\frac{\partial}{\partial \vec{x}'} f(|\vec{x} - \vec{x}'|)$$

Radiation From Relativistic Electrons



$$\phi(\vec{x}, t) = \frac{1}{8\pi^2 \epsilon_0} \int d^3x' dt' d\omega \frac{\rho(\vec{x}', t')}{|\vec{x} - \vec{x}'|} e^{i\omega[|\vec{x} - \vec{x}'|/c - (t - t')]}$$

$$\vec{A}(\vec{x}, t) = \frac{\mu_0}{8\pi^2} \int d^3x' dt' d\omega \frac{\vec{J}(\vec{x}', t')}{|\vec{x} - \vec{x}'|} e^{i\omega[|\vec{x} - \vec{x}'|/c - (t - t')]}$$

$$\rho(\vec{x}, t) = q\delta^3(\vec{x} - \vec{r}(t)) \quad \vec{J}(\vec{x}, t) = q\vec{v}(t)\delta^3(\vec{x} - \vec{r}(t))$$

$$\phi(\vec{x}, t) = \frac{q}{8\pi^2 \epsilon_0} \int dt' d\omega \frac{1}{|\vec{x} - \vec{r}(t')|} e^{i\omega[|\vec{x} - \vec{r}(t')|/c - (t - t')]}$$

$$\vec{A}(\vec{x}, t) = \frac{q\mu_0}{8\pi^2} \int dt' d\omega \frac{\vec{v}(t')}{|\vec{x} - \vec{r}(t')|} e^{i\omega[|\vec{x} - \vec{r}(t')|/c - (t - t')]}$$

Lienard-Weichert Potentials



$$\phi(\vec{x}, t) = \frac{q}{4\pi\epsilon_0} \int dt' \frac{\delta\left(|\vec{x} - \vec{r}(t')|/c - (t - t')\right)}{|\vec{x} - \vec{r}(t')|}$$

$$\vec{A}(\vec{x}, t) = \frac{q\mu_0}{4\pi} \int dt' \frac{\vec{v}(t') \delta\left(|\vec{x} - \vec{r}(t')|/c - (t - t')\right)}{|\vec{x} - \vec{r}(t')|}$$

$$\phi(\vec{x}, t) = \frac{q}{4\pi\epsilon_0} \left(\frac{1}{|\vec{x} - \vec{r}(t')| (1 - \hat{n} \cdot \vec{\beta}(t'))} \right)_{ret}$$

$$\vec{A}(\vec{x}, t) = \frac{q\mu_0}{4\pi} \left(\frac{\vec{v}(t')}{|\vec{x} - \vec{r}(t')| (1 - \hat{n} \cdot \vec{\beta}(t'))} \right)_{ret}$$

EM Field Radiated



$$\phi(\vec{x}, t) = \frac{q}{8\pi^2 \epsilon_0} \int dt' d\omega \frac{1}{|\vec{x} - \vec{r}(t')|} e^{i\omega[|\vec{x} - \vec{r}(t')|/c - (t-t')]}$$

$$\vec{A}(\vec{x}, t) = \frac{q\mu_0}{8\pi^2} \int dt' d\omega \frac{\vec{v}(t')}{|\vec{x} - \vec{r}(t')|} e^{i\omega[|\vec{x} - \vec{r}(t')|/c - (t-t')]}$$

$$\vec{E} = -\vec{\nabla}\phi - \frac{\partial\vec{A}}{\partial t} \quad \vec{B} = \vec{\nabla} \times \vec{A}$$

$$\vec{E} = \frac{q}{4\pi\epsilon_0} \left[\frac{\hat{n} - \vec{\beta}}{\gamma^2 (1 - \hat{n} \cdot \vec{\beta})^3 R^2} \right]_{ret} + \frac{q}{4\pi\epsilon_0 c} \left[\frac{\hat{n} \times \left\{ (\hat{n} - \vec{\beta}) \times \dot{\vec{\beta}} \right\}}{(1 - \hat{n} \cdot \vec{\beta})^3 R} \right]_{ret}$$

$$\vec{B} = \hat{n} \times \vec{E} / c$$

Velocity Field

Acceleration Field



$$\vec{\nabla} \frac{1}{|\vec{x} - \vec{r}(t')|} = - \frac{\hat{n}}{|\vec{x} - \vec{r}(t')|^2} \quad \vec{\nabla} |\vec{x} - \vec{r}(t')| = \hat{n}$$

$$\frac{d}{dt'} \frac{1}{|\vec{x} - \vec{r}(t')|} = \frac{\hat{n} \cdot \vec{\beta} c}{|\vec{x} - \vec{r}(t')|^2} \quad \frac{d\hat{n}}{dt'} = \frac{-d\vec{r}/dt' + \hat{n}(\hat{n} \cdot d\vec{r}/dt')}{|\vec{x} - \vec{r}(t')|} \quad \dots$$

$$-\nabla \phi(\vec{x}, t) = \frac{q}{8\pi^2 \epsilon_0} \int dt' d\omega \frac{\hat{n} (1 - i\omega |\vec{x} - \vec{r}(t')|/c)}{|\vec{x} - \vec{r}(t')|^2} e^{i\omega [|\vec{x} - \vec{r}(t')|/c - (t-t')]}$$

$$-\frac{\partial}{\partial t} \vec{A}(\vec{x}, t) = \frac{q\mu_0}{8\pi^2} \int dt' d\omega \frac{\vec{v}(t') i\omega}{|\vec{x} - \vec{r}(t')|} e^{i\omega [|\vec{x} - \vec{r}(t')|/c - (t-t')]}$$

$$\frac{d}{dt'} e^{i\omega [|\vec{x} - \vec{r}(t')|/c - (t-t')]} = i\omega (1 - \vec{\beta}(t') \cdot \hat{n}(t')) e^{i\omega [|\vec{x} - \vec{r}(t')|/c - (t-t')]}$$



$$\vec{E}(\vec{x}, t) = \frac{q}{8\pi^2 \epsilon_0} \int dt' d\omega e^{i\omega[|\vec{x}-\vec{r}(t')|/c-(t-t')]} \left[\frac{\hat{n}}{|\vec{x}-\vec{r}(t')|^2} + \frac{i\omega(\vec{\beta}-\hat{n})}{c|\vec{x}-\vec{r}(t')|} \right]$$

integrate by parts to get final result

$$\vec{E}(\vec{x}, t)_{vel} = \frac{q}{8\pi^2 \epsilon_0} \int dt' d\omega \frac{e^{i\omega[|\vec{x}-\vec{r}(t')|/c-(t-t')]}}{(1-\vec{\beta} \cdot \hat{n})^2 |\vec{x}-\vec{r}(t')|^2} \times \left[\begin{array}{l} \hat{n} \left(1 - 2\vec{\beta} \cdot \hat{n} + (\vec{\beta} \cdot \hat{n})^2 + \vec{\beta} \cdot \hat{n} - (\vec{\beta} \cdot \hat{n})^2 + \vec{\beta} \cdot \hat{n} - (\vec{\beta} \cdot \hat{n})^2 \right) \\ -\beta^2 + (\vec{\beta} \cdot \hat{n})^2 \\ -\vec{\beta} \left(1 - \vec{\beta} \cdot \hat{n} + \vec{\beta} \cdot \hat{n} - (\vec{\beta} \cdot \hat{n})^2 - \beta^2 + (\vec{\beta} \cdot \hat{n})^2 \right) \end{array} \right]$$



$$\begin{aligned}\vec{E}(\vec{x}, t)_{acc} &= \frac{q}{8\pi^2 \epsilon_0 c} \int dt' d\omega \frac{e^{i\omega[|\vec{x}-\vec{r}(t')|/c-(t-t')]}}{(1-\vec{\beta} \cdot \hat{n})^2 |\vec{x}-\vec{r}(t')|} \\ &\times \begin{bmatrix} \hat{n}(\dot{\vec{\beta}} \cdot \hat{n}) \\ -\dot{\vec{\beta}}(1-\vec{\beta} \cdot \hat{n}) - \vec{\beta}(\dot{\vec{\beta}} \cdot \hat{n}) \end{bmatrix} \\ &= \frac{q}{8\pi^2 \epsilon_0 c} \int dt' d\omega \frac{e^{i\omega[|\vec{x}-\vec{r}(t')|/c-(t-t')]}}{(1-\vec{\beta} \cdot \hat{n})^2 |\vec{x}-\vec{r}(t')|} \left[\hat{n} \times \left\{ (\hat{n} - \vec{\beta}) \times \dot{\vec{\beta}} \right\} \right]\end{aligned}$$

Larmor's Formula Verified



For small velocities can neglect retardation

$$\vec{E}(\vec{x}, t)_{acc} = \frac{q}{4\pi\epsilon_0 c} \left[\hat{n} \times \left\{ \hat{n} \times \dot{\vec{\beta}} \right\} \right] / R$$

$$\frac{dP}{d\Omega} = \frac{q^2}{16\pi^2 \epsilon_0^2 \mu_0 c^3} \left| \left[\hat{n} \times \left\{ \hat{n} \times \dot{\vec{\beta}} \right\} \right] \right|^2$$

$$= \frac{q^2}{16\pi^2 \epsilon_0 c^3} |\dot{\vec{v}}|^2 \sin^2 \Theta$$

$$P = \frac{q^2}{6\pi\epsilon_0 c^3} |\dot{\vec{v}}|^2$$

⊖ Angle between acceleration
and propagation directions

Classical Dipole
Radiation Pattern

Relativistic Peaking



In far field after short acceleration

$$\frac{dP(t')}{d\Omega} = \frac{q^2}{16\pi^2 \epsilon_0 c} \frac{\left| \hat{n} \times \left\{ (\hat{n} - \vec{\beta}) \times \dot{\vec{\beta}} \right\} \right|^2}{(1 - \hat{n} \cdot \vec{\beta})^5}$$

$$\frac{dP(t')}{d\Omega} = \frac{q^2 \dot{\beta}^2}{16\pi^2 \epsilon_0 c} \frac{\sin^2 \theta}{(1 - \beta \cos \theta)^5}$$

$$\theta_{\max} \rightarrow \frac{1}{2\gamma}$$

For circular motions

$$\frac{dP(t')}{d\Omega} = \frac{q^2}{16\pi^2 \epsilon_0 c} \frac{\dot{\beta}^2}{(1 - \beta \cos \theta)^3} \left[1 - \frac{\sin^2 \theta \cos^2 \varphi}{\gamma^2 (1 - \beta \cos \theta)^2} \right]$$

Spectrum Radiated by Motion



$$\frac{dE}{d\Omega} = \int_{-\infty}^{\infty} \frac{dP}{d\Omega} dt = \int_{-\infty}^{\infty} \vec{E} \times \vec{H} \cdot \hat{n} R^2 dt = \frac{1}{c\mu_0} \int_{-\infty}^{\infty} (\vec{E} \cdot \vec{E}) R^2 dt =$$

$$\frac{1}{c\mu_0} \left(\frac{q}{8\pi^2 \epsilon_0 c} \right)^2 \int_{-\infty}^{\infty} \int \int \int \int \left[\frac{\hat{n} \times \left\{ (\hat{n} - \vec{\beta}) \times \dot{\vec{\beta}} \right\}}{(1 - \hat{n} \cdot \vec{\beta})^2} (t') \right] \cdot \left[\frac{\hat{n} \times \left\{ (\hat{n} - \vec{\beta}) \times \dot{\vec{\beta}} \right\}}{(1 - \hat{n} \cdot \vec{\beta})^2} (t'') \right] \times$$

$$e^{i\omega \left[R\sqrt{1-2\hat{n}\cdot\vec{r}(t')/R+(\hat{n}\cdot\vec{r}(t'))^2/R^2}/c-t+t' \right]} e^{i\omega' \left[R\sqrt{1-2\hat{n}\cdot\vec{r}(t'')/R+(\hat{n}\cdot\vec{r}(t''))^2/R^2}/c-t+t'' \right]} dt' d\omega dt'' d\omega' dt =$$

clearing the unprimed time integral and omega prime
integral with delta representation

$$\frac{2\pi}{c\mu_0} \left(\frac{q}{8\pi^2 \epsilon_0 c} \right)^2 \int_{-\infty}^{\infty} \int \int \left[\frac{\hat{n} \times \left\{ (\hat{n} - \vec{\beta}) \times \dot{\vec{\beta}} \right\}}{(1 - \hat{n} \cdot \vec{\beta})^2} (t') \right] \cdot \left[\frac{\hat{n} \times \left\{ (\hat{n} - \vec{\beta}) \times \dot{\vec{\beta}} \right\}}{(1 - \hat{n} \cdot \vec{\beta})^2} (t'') \right]$$

$$\times e^{i\omega \left[-\hat{n}\cdot\vec{r}(t')/c+t' \right]} e^{-i\omega \left[-\hat{n}\cdot\vec{r}(t'')/c+t'' \right]} dt' dt'' d\omega$$



$$\frac{d^2 E}{d\omega d\Omega} = \frac{q^2}{32\pi^3 \epsilon_0 c} \left| \int_{-\infty}^{\infty} \frac{\hat{n} \times \left\{ (\hat{n} - \vec{\beta}) \times \dot{\vec{\beta}} \right\}}{(1 - \hat{n} \cdot \vec{\beta})^2} e^{i\omega[-\hat{n} \cdot \vec{r}(t')/c - t + t']} dt' \right|^2$$

$$\frac{d^2 E}{d\omega d\Omega} = \frac{q^2 \omega^2}{32\pi^3 \epsilon_0 c} \left| \int_{-\infty}^{\infty} \hat{n} \times (\hat{n} \times \vec{\beta}) e^{i\omega[t' - \hat{n} \cdot \vec{r}(t')/c]} dt' \right|^2$$

Factor of two difference from Jackson because he combines positive frequency and negative frequency contributions in one positive frequency integral. I don't like because Parseval's formula, etc. don't work! We'll perform this calculation in new intensity regimes of pulsed lasers.